

In the claims:

1-4. (Cancelled)

5. **(Currently Amended)** A composition comprising a cyclodextrin-containing polymer and a therapeutic agent and a complexing agent comprising at least one functional group and at least one host/guest moiety that forms an inclusion complex with a host/guest moiety of said polymer, wherein the polymer, the therapeutic agent, and the complexing agent are separate molecules.


6. **(Previously Presented)** A composition of claim 5, wherein said therapeutic agent is selected from an antibiotic, a steroid, a polynucleotide, small molecule pharmaceutical, a virus, a plasmid, a peptide, a peptide fragment, a chelating agent, a biologically active macromolecule, and mixtures thereof.

7. **(Original)** A composition of claim 6, wherein said therapeutic agent is a polynucleotide.

8-10. (Cancelled)


11. **(Previously Presented)** A composition of claim 5, wherein the host/guest of the complexing agent is selected from adamantyl, naphthyl, cholesterol, cyclodextrin, and mixtures thereof.

12. **(Previously Presented)** A composition of claim 5, wherein the complexing agent is a compound of the formula:



wherein

J is $-\text{NH}-$, $-\text{C}(=\text{O})\text{NH}-\text{CH}_2$, $-\text{NH}-\text{C}(=\text{O})-(\text{CH}_2)_d-$, $-\text{CH}_2\text{SS}-$, $-\text{C}(=\text{O})\text{O}-(\text{CH}_2)_e-\text{O}-\text{P}(=\text{O})(\text{O}-$



Y is an additional host-guest functionality;

R^1 is $-(\text{CH}_2)-\text{CO}_2\text{H}$, an ester or salt thereof; or $-(\text{CH}_2)_a-\text{CONH}_2$;

PEG is $-\text{O}(\text{CH}_2\text{CH}_2\text{O})_z-$, where z varies from 2 to 500;

L is H, $-\text{NH}$, $-\text{NH}-(\text{C}=\text{O})-(\text{CH}_2)_e-(\text{C}=\text{O})-\text{CH}_2-$, $-\text{S}(=\text{O})_2-\text{HC}=\text{CH}-$, $-\text{SS}-$, $-\text{C}(=\text{O})\text{O}-$, or a carbohydrate residue;

a is 0 or 1;

b is 0 or 1;

d ranges from 0 to 6;

e ranges from 1 to 6;

n ranges from 0 to 6;


q ranges from 1 to 5;

w ranges from 1 to 5;

y is 1; and


x is 0 or 1.

13. **(Previously Presented)** A composition of claim 5, wherein the complexing agent is a compound of the formula:



wherein

J is $-\text{NH}-$, $-\text{C}(=\text{O})\text{NH}-(\text{CH}_2)_d-$, $-\text{NH}-\text{C}(=\text{O})-(\text{CH}_2)_d-$, $-\text{CH}_2\text{SS}-$, $-\text{C}(=\text{O})\text{O}-(\text{CH}_2)_e-\text{O}-\text{P}(=\text{O})(\text{O}-$



Y is an additional host-guest functionality;

R^1 is $-(\text{CH}_2)-\text{CO}_2\text{H}$, an ester or salt thereof; or $-(\text{CH}_2)_a-\text{CONH}_2$;

PEG is $-\text{O}(\text{CH}_2\text{CH}_2\text{O})_z-$, where z varies from 2 to 500;

L is H, $-\text{NH}-$, $-\text{NH}-(\text{C}=\text{O})-(\text{CH}_2)_e-(\text{C}=\text{O})-\text{CH}_2-$, $-\text{S}(=\text{O})_2-\text{HC}=\text{CH}-$, $-\text{SS}-$, $-\text{C}(=\text{O})\text{O}-$, or a carbohydrate residue;

a is 0 or 1;

b is 0 or 1;

d ranges from 0 to 6;

e ranges from 1 to 6;

n ranges from 0 to 6;

y is 1; and

x is 0 or 1.

14. **(Previously Presented)** A composition of claim 5, wherein the at least one functional group includes a group selected from a ligand, a nuclear localization signal, an endosomal release peptide, an endosomal release polymer, or a membrane permeabilization agent.

15. **(Previously Presented)** A composition of claim 5, wherein the at least one functional group includes a moiety that increases the solubility of the composition under biological conditions relative to a composition of the polymer and therapeutic agent alone.
16. **(Previously Presented)** A composition of claim 5, wherein the at least one functional group includes a moiety that stabilizes the composition under biological conditions relative to a composition of the polymer and therapeutic agent alone.
17. **(Previously Presented)** A composition of claim 5, wherein the at least one functional group includes a therapeutic agent reversibly bound to the complexing agent.
18. **(Previously Presented)** A composition of claim 5, wherein the complexing agent further comprises a spacer group.
19. **(Previously Presented)** A composition of claim 5, wherein the polymer comprises a host moiety that forms an inclusion complex with a guest moiety of the complexing agent.
20. **(Previously Presented)** A composition of claim 5, wherein the polymer comprises a guest moiety that forms an inclusion complex with a host moiety of the complexing agent.
21. **(Previously Presented)** A composition of claim 20, wherein the guest moiety is an adamantyl group and the host moiety is a cyclodextrin moiety.